Acer x freemanii AUTUMN BLAZE Jeffersred Freeman's Maple



SEASONAL COLOURS

| jan | feb | mar | apr | mei | jun | jul | aug | sep | okt | nov | dec |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|

TYPES OF PLANTING

Tree types: standard trees, feathered trees, multi-stemmed trees, climate trees, shade trees | Topiary on stem: block, multi-stem umbrella

USE

Location: street, avenue, square, park, central reservation, in containers, roof garden, large garden, cemetery, traffic areas, industrial zones | **Pavement**: none, open, sealed | **Planting concepts**: Climate planting, urban planting

CHARACTERISTICS

Crown shape: wide egg-shaped | Crown structure: dense | Height: 12 - 17 m | Width: 9 - 12 m | Winter hardiness zone: 4A - 9B

ASPECTS

Wind: tolerant to wind | **Soil**: loess, sabulous clay, peaty, heavy clay, light clay, sand, loamy soil, all soils | **Nutrient level**: moderately rich in nutrients, rich in nutrients | **Soil moisture level**: dry, moist, wet | **Light requirements**: sun, partial shade | **pH range**: acidic, neutral, alkaline | **Host plant/forage plant**: bees, butterflies | **Extreme environments**: tolerant to dryness, tolerates air pollution, tolerates heat

PLANTKENMERKEN

Flowers: corymbose, discrete | Flower colour: yellow-green | Flowering period: March - April | Leaf colour: buds bronze, bright green, underside grey-green | Leaves: deciduous, palmate, thick, opposite, doubly serrate, lobate | Autumn colour: orange-red, scarlet | Fruits: rarely fruits, winged | Fruit colour: yellow-brown, green | Bark colour: grey-brown | Bark: furrowed | Twig colour: red-brown | Twigs: striped, with lenticels | Root system: deep, shallow, fine roots



TreeEbb