Tilia americana 'Redmond' American linden 'Redmond', American basswood 'Redmond'



SEASONAL COLOURS

| jan | feb | mar | apr | mei | jun | jul | aug | sep | okt | nov | dec |
|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|-----|

TYPES OF PLANTING

Tree types: standard trees, shade trees | Topiary on stem: block, candelabra

USE

Location: street, avenue, square, park, central reservation, large garden, cemetery, countryside, ecological zone, windbreak | **Pavement**: none, open, sealed | **Planting concepts**: Eco planting, Landscape planting, shade-tolerant, urban planting

CHARACTERISTICS

Crown shape: ovoid, pyramidal | Crown structure: dense | Height: 20 - 25 m | Width: 10 - 12 m | Winter hardiness zone: 5B - 8B

ASPECTS

Wind: tolerant to wind | Soil: loess, sabulous clay, peaty, heavy clay, light clay, sand, loamy soil, all soils | Nutrient level: moderately rich in nutrients, rich in nutrients | Soil moisture level: dry, moist | Light requirements: sun, partial shade, shade | pH range: acidic, neutral, alkaline | Host plant/forage plant: bees, butterflies, nectar value 5, pollen value 5 | Extreme environments: tolerant to dryness, tolerates air pollution, limited to rare infestation by lice

PLANTKENMERKEN

Flowers: corymbose, striking, pendulous, flower-scented | Flower colour: pale yellow | Flowering period: June - June | Leaf colour: pale green | Leaves: deciduous, cordate, big, polished, hairy, serrate | Autumn colour: yellow | Fruits: hairy, discrete, nut | Fruit colour: grey | Bark colour: grey | Bark: slightly furrowed | Twig colour: olive green | Twigs: bare, vertical branches, thickly branched | Root system: deep, extensive, coarse roots



